

Towards supersymmetric cosmology in M–theory

Krzysztof A. Meissner and Marek Olechowski

Institute of Theoretical Physics

Warsaw University

Hoża 69, 00-681 Warsaw, Poland

Abstract

We present a new solution in the heterotic M–theory in which the metric depends on (cosmic) time. The solution preserves $N = 1$ supersymmetry in 4 dimensions in the leading order of the $\kappa^{2/3}$ expansion. It is the first example of the time–dependent supersymmetric solution in M–theory on $S^1/\mathbb{Z}_2$. It describes expanding 4–dimensional space–time with shrinking orientifold interval and static Calabi–Yau internal space.

The M-theoretic extension of the heterotic $E_8 \times E_8$ string leads to a geometric picture of two 10-dimensional walls (branes) at the ends of a finite interval along eleventh dimension [1, 2]. While the supergravity multiplet (graviton, gravitino and antisymmetric tensor fields) can penetrate in the $d = 11$ bulk, the two E_8 gauge multiplets are confined to the two walls, respectively. The role of the walls can in general be played by higher dimensional p-branes that support gauge groups. When six of the dimensions are compactified (at the lowest order on a Calabi–Yau space) one obtains phenomenologically promising models with gauge fields living on the $d = 4$ walls. One should however remember that such a compactification should be consistent with the 11-dimensional equations of motion.

Within this heterotic M-theory it is possible to unify the gravitational constant with the gauge coupling constants at the M-theory fundamental scale. Assuming such a unification and using measured values of the Newton’s constant G_N and of the gauge coupling constants one can deduce scales of compactification. These scales of the internal dimensions are of the order of 10^{15} GeV for the x^{11} interval and 10^{16} GeV for the Calabi–Yau space [2, 3, 4]. These energy scales are very similar to the characteristic scales in the Early Universe, for example during inflationary period. Therefore it is natural to ask whether cosmological solutions (with the metric depending on time) are possible in M-theory.

The problem of finding cosmological solutions in M-theory has been addressed in several papers. Some of them discuss solutions within strongly coupled limit of the heterotic string theory (heterotic M-theory). One of the approaches in this case consists in using an effective 5-dimensional theory with walls [5]. The domain wall solutions [6] are generalized by introducing time dependence of their parameters. This approach can be justified [7] because the scale of the x^{11} interval is lower than the Calabi–Yau scale. But one should check *a posteriori* whether the 11-dimensional equations of motion are satisfied and it is a very stringent requirement. In another approach one looks for solutions in the theory with walls by continuation of p-branes to the Euclidean space [8].

Other papers discuss solutions within 11- or 10-dimensional supergravities without walls (general M-theory, less promising phenomenologically). There one assumes different compact, internal manifolds (for example squashed spheres [9] or tori [10]) or compactification by the Scherk–Schwarz mechanism [11] (giving a static Universe with domain–walls).

In the case of the heterotic M–theory none of the cosmological solutions proposed up to now preserves $N = 1$ $d = 4$ supersymmetry. The purpose of this paper is to generalize the result of ref. [2] and prove that such a supersymmetric solution is possible.

The low energy limit of the heterotic M–theory is given by the following lagrangian [1]:

$$\begin{aligned}
\mathcal{L} = & \frac{1}{\kappa^2} \int_{M^{11}} d^{11}x \sqrt{g} \left[-\frac{1}{2}R - \frac{1}{2} \bar{\psi}_I \Gamma^{IJK} D_J \left(\frac{\Omega + \hat{\Omega}}{2} \right) \psi_K - \frac{1}{48} G_{IJKL} G^{IJKL} \right. \\
& - \frac{\sqrt{2}}{384} \left(\bar{\psi}_I \Gamma^{IJKLMN} \psi_N + 12 \bar{\psi}^J \Gamma^{KL} \psi^M \right) (G_{JKLM} + \hat{G}_{JKLM}) \\
& \left. - \frac{\sqrt{2}}{3456} \epsilon^{I_1 I_2 \dots I_{11}} C_{I_1 I_2 I_3} G_{I_4 \dots I_7} G_{I_8 \dots I_{11}} \right] \\
& + \frac{1}{4\pi(4\pi\kappa^2)^{2/3}} \sum_{n=1}^2 \int_{M_n^{10}} d^{10}x \sqrt{g} \left[-\frac{1}{4} F_{nXY}^a F_n^{aXY} - \frac{1}{2} \bar{\chi}_n^a \Gamma^X D_X (\hat{\Omega}) \chi_n^a \right. \\
& \left. - \frac{1}{8} \bar{\psi}_X \Gamma^{YZ} \Gamma^X (F_{nYZ}^a + \hat{F}_{nYZ}^a) \chi_n^a + \frac{\sqrt{2}}{48} (\bar{\chi}_n^a \Gamma^{XYZ} \chi_n^a) \hat{G}_{XYZ11} \right]
\end{aligned} \tag{1}$$

where $I, J, K, \dots = 1, 2, \dots, 11$; $X, Y, Z \dots = 1, 2, \dots, 10$ and $n = 1, 2$ counts the 10–dimensional boundaries (walls) of the space. The first integral in (1) describes the supergravity in the 11–dimensional bulk (metric g_{IJ} , three-form C_{IJK} and spin 3/2 gravitino ψ_I) while the second one describes interactions of the super Yang–Mills fields (gauge field strengths F_n^a and gauginos χ_n^a) living on two 10–dimensional walls. The field strength of C is given by $G_{IJKL} = (\partial_I C_{JKL} \pm \text{permutations})$ and $\hat{G}$ is the supercovariant field strength. The signature of the space-time manifold M^{11} is $(- + \dots +)$; gamma matrices obey $\{\Gamma_I, \Gamma_J\} = 2g_{IJ}$ and $\Gamma_{I_1 I_2 \dots I_n} = (1/n!) (\Gamma_{I_1} \Gamma_{I_2} \dots \Gamma_{I_n} \pm \text{permutations})$.

In the above lagrangian only the first two terms in the long wavelength expansion are kept. They are of relative order $\kappa^{2/3}$. All higher order terms (order $\kappa^{4/3}$ or higher) will be consistently dropped in this paper.

We now try to find a time–dependent solution that preserves at least some of the supersymmetries generalizing the method employed in [2].

The supersymmetry transformation law for ψ_I is

$$\delta\psi_I = D_I \epsilon + \frac{\sqrt{2}}{288} (\Gamma_{IJKLM} - 8g_{IJ} \Gamma_{KLM}) \hat{G}^{JKLM} \epsilon. \tag{2}$$

The condition for a spinor ϵ to generate an unbroken supersymmetry is the van-

ishing of the r.h.s. of (2):

$$D_I \epsilon + \frac{\sqrt{2}}{288} (\Gamma_{IJKLM} - 8g_{IJ}\Gamma_{KLM}) G^{JKLM} \epsilon = 0. \quad (3)$$

In the above equation we use the covariant (and not supercovariant) field strength because we assume that vacuum expectation values of all fermion fields vanish.

To the zeroth order in $\kappa^{2/3}$ the 11-dimensional space-time in the heterotic M-theory is of the form $M^{11} = M^4 \times X^6 \times S^1/\mathbb{Z}_2$. Indices $A, B, C, \dots$ run from 5 to 10 and are tangent to X^6 ; $a, b, c, \dots = 1, 2, 3$ are holomorphic indices tangent to X^6 , and $\bar{a}, \bar{b}, \bar{c}, \dots = 1, 2, 3$ are analogous antiholomorphic indices; $\mu, \nu, \rho, \dots = 1, 2, 3, 4$ are $d = 4$ space-time indices while $i, j, k, \dots = 2, 3, 4$ are space indices.

We are looking for a solution of the supersymmetry condition (3) in the presence of nonvanishing vev of G_{JKLM} of the order of $\kappa^{2/3}$. First we perturb the metric:

$$ds^2 = (\eta_{\mu\nu} + b_{\mu\nu}) dx^\mu dx^\nu + 2(g_{a\bar{b}} + h_{a\bar{b}}) dx^a dx^{\bar{b}} + (1 + c)(dx^{11})^2 \quad (4)$$

where b, h , and c are functions of order $\kappa^{2/3}$ and we will neglect higher order corrections. We take the zeroth order metric on M^4 to be the Minkowski one $\eta_{\mu\nu} dx^\mu dx^\nu$. X^6 is to zeroth order a Calabi-Yau manifold with a metric g_{AB} with non-zero components $g_{a\bar{b}} = g_{\bar{b}a}$. A Kähler form on X^6 is ω_{AB} with non-zero components $\omega_{a\bar{b}} = -i g_{a\bar{b}} = -\omega_{\bar{b}a}$.

Under a perturbation of the metric $g_{IJ} \rightarrow g_{IJ} + \delta g_{IJ}$, to first order in δg , the covariant derivative of a spinor changes by $D_I \epsilon \rightarrow D_I \epsilon - \frac{1}{4} D_J \delta g_{KI} \Gamma^{JK} \epsilon$. Introducing the covariantly constant (with respect to the unperturbed metric) spinor field ϵ_0 we write

$$\epsilon = e^p \epsilon_0, \quad (5)$$

p being also of order $\kappa^{2/3}$.

Then the first term in eq. (3) is

$$\begin{aligned} dx^I D_I \epsilon = \frac{1}{4} \big\{ & 4dx^I \partial_I p + dx^\mu \left(\partial_{11} b_{\mu\rho} \Gamma^\rho + \partial_{\bar{b}} b_{\mu\rho} \Gamma^{\rho\bar{b}} - D_\rho b_{\sigma\mu} \Gamma^{\rho\sigma} \right) \\ & + dx^a \left(-g^{\bar{b}\bar{c}} D_{\bar{b}} h_{a\bar{c}} + \partial_{11} h_{a\bar{b}} \Gamma^{\bar{b}} - D_{\bar{b}} h_{a\bar{c}} \Gamma^{\bar{b}\bar{c}} - \partial_\mu h_{a\bar{b}} \Gamma^{\mu\bar{b}} \right) \\ & + dx^{\bar{b}} \left(g^{a\bar{c}} D_{\bar{c}} h_{a\bar{b}} \right) + dx^{11} \left(-\partial_{\bar{b}} c \Gamma^{\bar{b}} - \partial_\mu c \Gamma^\mu \right) \big\} \epsilon. \end{aligned} \quad (6)$$

The $\mathbb{Z}_2$ invariance requires that the supersymmetry transformation parameter ϵ obeys chirality condition $\Gamma^{11} \epsilon = \epsilon$. To simplify calculations it is also convenient

to assume that $\Gamma^a \epsilon = 0$. For spinors which are singlets of the $SU(3)$ holonomy group of X^6 this reduces to the Weyl condition $\gamma^5 \epsilon = -\epsilon$ in $d = 4$ (where $\gamma^5 = -i\Gamma^1\Gamma^2\Gamma^3\Gamma^4$ and condition $\Gamma^{\bar{a}}\epsilon = 0$ would correspond to the opposite 4-dimensional chirality). From Weyl-type solutions of the supersymmetry condition we can construct Majorana-type solutions if needed.

Supersymmetry condition (3) with the perturbed metric (4) can be solved only if some of the components of the field strength tensor G are nonzero. It was shown in [2] that the following components of G

$$\alpha = \omega^{AB}\omega^{CD}G_{ABCD}, \quad (7)$$

$$\beta_A = \omega^{BC}G_{ABC11}, \quad (8)$$

$$\theta_{AB} = \omega^{CD}G_{ABCD} \quad (9)$$

must be nonzero in the presence of sources. Looking for solutions depending on space-time coordinates we have to assume that there are more nonvanishing components of G . If we are looking for supersymmetric solutions the only additional components allowed are of the type $G_{\mu11a\bar{b}}$. Therefore, we define

$$\zeta_\mu = i\omega^{AB}G_{\mu11AB}. \quad (10)$$

There are no sources entering the Bianchi identity for this quantity. It was argued in [2] that "sourceless" components of G should vanish since G with indices tangent to the internal space must be cohomologically trivial. This argument does not apply to $G_{\mu11a\bar{b}}$ because $d = 4$ space-time is not compact and boundaryless so even without sources allows for constant field strengths.

With the above definitions the second term in the supersymmetry condition (3) can be written as

$$\begin{aligned} & \frac{\sqrt{2}dx^I}{288} (\Gamma_{IJKLM} - 8g_{IJ}\Gamma_{KLM}) G^{JKLM} \epsilon = \\ & = \frac{\sqrt{2}}{288} \left\{ dx^\mu \left(24\zeta_\mu - 3\alpha\Gamma_\mu - 12i\eta_{\mu\rho}\beta_{\bar{b}}\Gamma^{\rho\bar{b}} - 12\eta_{\mu\rho}\zeta_\sigma\Gamma^{\rho\sigma} \right) \right. \\ & \quad + dx^a \left(36i\beta_a + (36i\theta_{a\bar{b}} - 3\alpha g_{a\bar{b}})\Gamma^{\bar{b}} + (36G_{11a\bar{b}\bar{c}} + 12ig_{a\bar{c}}\beta_{\bar{b}})\Gamma^{\bar{b}\bar{c}} \right. \\ & \quad \left. \left. + (72G_{\mu11a\bar{b}} + 12g_{a\bar{b}}\zeta_\mu)\Gamma^{\mu\bar{b}} \right) \right. \\ & \quad \left. + dx^{\bar{b}} (12i\beta_{\bar{b}}) + dx^{11} \left(-3\alpha - 24i\beta_{\bar{b}}\Gamma^{\bar{b}} - 24\zeta_\mu\Gamma^\mu \right) \right\} \epsilon. \quad (11) \end{aligned}$$

Substituting (6) and (11) into (3) we obtain the following equations for the perturbations of the metric and the supersymmetry generating spinor:

$$D_\rho b_{\sigma\mu} - D_\sigma b_{\rho\mu} = \frac{\sqrt{2}}{6} (g_{\mu\sigma}\zeta_\rho - g_{\mu\rho}\zeta_\sigma), \quad (12)$$

$$\partial_\mu h_{a\bar{b}} = \frac{\sqrt{2}}{6} g_{a\bar{b}} \zeta_\mu + \sqrt{2} G_{\mu 11 a \bar{b}}, \quad (13)$$

$$\partial_\mu c = -\frac{\sqrt{2}}{3} \zeta_\mu, \quad (14)$$

$$\partial_\mu p = -\frac{\sqrt{2}}{12} \zeta_\mu, \quad (15)$$

$$\partial_{\bar{a}} b_{\mu\nu} = \frac{\sqrt{2}}{6} i \beta_{\bar{a}} \eta_{\mu\nu}, \quad (16)$$

$$\partial_{11} b_{\mu\nu} = \frac{\sqrt{2}}{24} \alpha \eta_{\mu\nu}. \quad (17)$$

The other equations are the same as those obtained in [2].

One can check that eq. (3) has no solutions with nonzero components of G other than those given in eqs. (7–10).

In order to check the consistency of the above set of equations we use the following Bianchi identity

$$(dG)_{\mu\nu 11 a \bar{b}} = \partial_\mu G_{\nu 11 a \bar{b}} - \partial_\nu G_{\mu 11 a \bar{b}} = 0. \quad (18)$$

Contracting it with $\omega^{a\bar{b}}$ we get

$$\partial_\mu \zeta_\nu - \partial_\nu \zeta_\mu = 0. \quad (19)$$

It is easy to check that equations (18) and (19) are the integrability conditions for equations (12)–(15) (integrability conditions for (16) and (17) were discussed in [2]).

In the potential application of the solutions of (12)–(17) to cosmological situations the natural assumption is to allow for dependence of all fields on the cosmic time ($t = x^1$) but not on the space coordinates (we are perturbing flat Minkowski space–time so spaces with topologically different space-like sections are excluded). Therefore we assume

$$G_{i 11 a \bar{b}} = 0. \quad (20)$$

The equation of motion for G reads

$$\begin{aligned} D_I G^{IJKL} &= \frac{\sqrt{2}}{1152} \epsilon^{JKLI_4 \dots I_{11}} G_{I_4 I_5 I_6 I_7} G_{I_8 I_9 I_{10} I_{11}} \\ &\quad - \frac{1}{1728\pi} \left(\frac{\kappa}{4\pi} \right)^{2/3} \delta(x^{11}) \epsilon^{11JKLI_5 \dots I_{11}} \text{Tr} \left(AdA + \frac{2}{3} A^3 \right)_{I_5 I_6 I_7} G_{I_8 I_9 I_{10} I_{11}}. \end{aligned} \quad (21)$$

The r.h.s. of the above equation may be nonzero for general vacuum expectation values of G given by eqs. (7–10). However, as shown in ref. [2], quantities α , β_A and θ_{AB} are all of order $\kappa^{2/3}$. We are going to restrict our analysis to situations when ζ_μ is also of the order of $\kappa^{2/3}$ (ζ_μ of order 1 would not satisfy equations of motion at the leading order). For G satisfying this condition the r.h.s. of eq. (21) is of higher order ($\kappa^{4/3}$) and should be dropped.

From now on we will use for simplicity the cartesian coordinates to parametrize M^4 . Then, the equations of motion and Bianchi identities relevant for us are:

$$\partial_{11}G_{t11a\bar{b}} = 0, \quad (22)$$

$$-\partial_t G_{t11a\bar{b}} + D^c G_{c11a\bar{b}} + D^{\bar{c}} G_{\bar{c}11a\bar{b}} = 0, \quad (23)$$

$$(DG)_{t11a\bar{b}\bar{c}} = (DG)_{t11a\bar{b}\bar{c}} = (DG)_{ti11a\bar{b}} = 0. \quad (24)$$

By introducing $G_{t11a\bar{b}}$ we don't want to change to the lowest order the G components in the solution of [2] so we assume

$$\partial_t G_{t11a\bar{b}} = \partial_i G_{t11a\bar{b}} = \partial_c G_{t11a\bar{b}} = \partial_{\bar{c}} G_{t11a\bar{b}} = 0. \quad (25)$$

We can do this in a consistent way because the Bianchi identities for the added components of G ($G_{t11a\bar{b}}$) have no sources at the walls.

Contracting (25) with $\omega^{a\bar{b}}$ we get

$$\zeta_1 = \text{const}. \quad (26)$$

The explicit solution to eqs. (12)–(17) reads (we drop x^a and $x^{\bar{a}}$ dependence analyzed in [2]):

$$b_{tt}(t, x^{11}) = (\bar{\alpha}x^{11} + f_1(t)) \eta_{tt}, \quad (27)$$

$$b_{ij}(t, x^{11}) = (\bar{\alpha}x^{11} + \zeta t) \eta_{ij}, \quad (28)$$

$$c(t, x^{11}) = -2(f_2(x^{11}) + \zeta t), \quad (29)$$

$$p(t, x^{11}) = -\frac{1}{2}(\bar{\alpha}x^{11} + \zeta t). \quad (30)$$

where $\zeta = \frac{\sqrt{2}}{6}\zeta_1$, $\bar{\alpha}$ is an average value over the Calabi–Yau space of $\frac{\sqrt{2}}{24}\alpha$ and $f_1(t)$ and $f_2(x^{11})$ are arbitrary functions reflecting the freedom of coordinate choice of time and x^{11} . We have dropped the arbitrary constants to make the result look simpler.

It is important to analyse the volume of X^6 because it determines the value of the gauge coupling constants. Its dependence on the space-time coordinates is given by

$$\partial_\mu V_6 = \partial_\mu \int d^6x \sqrt{g^{(6)}} = \int d^6x \sqrt{g^{(6)}} g^{a\bar{b}} \partial_\mu h_{a\bar{b}}. \quad (31)$$

Substituting (13) into this formula we get

$$\begin{aligned} \partial_\mu V_6 &= \int d^6x \sqrt{g^{(6)}} g^{a\bar{b}} \left(\frac{\sqrt{2}}{6} g_{a\bar{b}} \zeta_\mu + \sqrt{2} G_{\mu 11 a \bar{b}} \right) \\ &= \int d^6x \sqrt{g^{(6)}} \left(\frac{\sqrt{2}}{2} \zeta_\mu + \sqrt{2} g^{a\bar{b}} G_{\mu 11 a \bar{b}} \right) = 0. \end{aligned} \quad (32)$$

In these equalities we used the definition of ζ_μ (eq. (10)) and the fact that the complex dimension of X^6 is equal to 3. Thus, V_6 depends only on x^{11} but not on time:

$$V_6(x^{11}) = V_6(0) - \frac{\sqrt{2}}{8} x^{11} \int d^6x \sqrt{g^{(6)}} \alpha. \quad (33)$$

This fact that the volume of Calabi–Yau space, V_6 , does not depend on time (as shown in eq. (32)) has important consequences. The squares of the gauge coupling constants are inversely proportional to the volume of the Calabi–Yau space V_6 . Therefore the gauge coupling constants do not depend explicitly on time.

On the other hand, the scale factor of the 3-dimensional space and the length of $S^1/\mathbb{Z}_2$ do change in time. To discuss possible expansion or contraction of internal or space-time dimensions it is convenient to fix $f_1(t)$. Apart from the x^{11} dependence, the usual cosmic time prescription amounts to putting $f_1(t) = 0$ and that is our choice (we also fix the x^{11} coordinate freedom by choosing $f_2(x^{11}) = 0$). It is interesting that when we choose the sign of ζ positive so that it describes expanding space (space Hubble parameter positive) then the solution describes shrinking x^{11} scale factor.

Since the length of the x^{11} interval in this solution may depend on time we have to reconsider the question of the connection between the Newton's constant and the gauge coupling constant at the GUT scale. For the time-independent solutions these relations read

$$G_N = \frac{\kappa_4^2}{8\pi} = \frac{\kappa^2}{8\pi R_{11} V_6}, \quad (34)$$

$$\alpha_{GUT} = \frac{(4\pi\kappa^2)^{2/3}}{V_6^0} \quad (35)$$

where $R_{11} = \pi\rho$ is the length of the $S^1/\mathbb{Z}_2$ interval, $V_6^0 = V_6(x^{11} = 0)$ is the volume of X^6 at our wall and $\bar{V}_6$ is the averaged (over x^{11}) volume of X^6 . Eliminating κ we get

$$R_{11} = \frac{(\alpha_{GUT} V_6^0)^{3/2}}{32\pi^2 G_N \bar{V}_6}. \quad (36)$$

Since in the linear approximation

$$\frac{1}{2}V_6^0 \leq \bar{V}_6 \leq V_6^0 \quad (37)$$

then knowing α_{GUT} and G_N and assuming $V_6^0 \approx M_{GUT}^{-6}$ we can calculate R_{11} – it turns out to be of the order

$$R_{11}^{-1} \approx 10^{15} \text{GeV}. \quad (38)$$

Plugging this value back into eq. (33) we have to require that the volume of X^6 at the other wall is positive which gives an upper bound on the length of $S^1/\mathbb{Z}_2$: $R_{11} < R_{11}^{crit}$. This critical length R_{11}^{crit} turns out to be close to the value (38) obtained from arguments based on phenomenology. One could expect that the evolution of the length of the x^{11} interval could violate the condition $R_{11} < R_{11}^{crit}$ for large enough negative t . However, this is not the case. R_{11}^{crit} is obtained from the condition that V_6 must be positive at both walls. We can see from eq. (33) that since x^{11} is the “comoving” coordinate the volume V_6 does not change with time and stays positive along the whole x^{11} interval. The physical length of the interval changes with time but this is “compensated” by the change of R_{11}^{crit} induced by the change of the Newton’s constant as shown in eq. (34).

Let us analyse the resulting metric in some more detail. We assume (as everywhere else in the paper) that we keep only linear terms in $\kappa^{2/3}$. Therefore the 5-dimensional metric reads

$$ds^2 = -\left(1 + \bar{\alpha}x^{11}\right) dt^2 + \left(1 + \bar{\alpha}x^{11} + \zeta t\right) (dx^i)^2 + (1 - 2\zeta t)(dx^{11})^2. \quad (39)$$

The Riemann tensor calculated from the above metric has an expansion in powers of $\bar{\alpha}$ and ζ starting from quadratic terms. Therefore the Riemann tensor components are of the order $\kappa^{4/3}$ and the equations of motion relating the Einstein tensor with the square of G are satisfied to order $\kappa^{2/3}$ (it is a similar situation as in [2]).

The metric (39) may depend on time while preserving 4 supersymmetry charges corresponding to $N = 1$ supersymmetry in $d = 4$ (this is similar to

the anti de Sitter space where supersymmetry can be preserved although the metric depends on the spatial coordinates). This fact is extremely important since preserved supersymmetry in a given background (with globally well defined fields) guarantees that this background satisfies the 11-dimensional equations of motion. Thus, the solution described in this paper is the first approximation (in the $\kappa^{2/3}$ expansion) to the time-dependent solutions in M-theory and it is the first example of such a solution.

Since 4-dimensional supersymmetry is preserved one should be able to identify an appropriate 4-dimensional gravitino field. This could be done using methods of ref. [12] generalized to the time dependent case.

The metric (39) is known in the linear approximation only, so we cannot make any definite statements about the global behaviour of the solution with time i.e. whether it is an inflationary-like solution or a power-like one or anything else. A similar problem exists already in the case of x^{11} dependence. There we also know the metric only in the linear approximation, but the problem with the global behaviour is less severe because the length of x^{11} interval is finite (although large).

One may note that the presence of nonzero $G_{t11a\bar{b}}$ may be important for the cosmological constant problem. It stems from the fact that the contribution of these components to vacuum energy has the opposite sign compared to the contribution of other components of G present in the static solution.

If we extend validity of the metric (39) to arbitrary time we can derive a rather surprising lower bound on the value of $G_{t11a\bar{b}}$. The argument goes as follows. One can see that the speed of light in the x^{11} direction is given by

$$v_{11}^2 = \frac{1 + \bar{\alpha}x^{11}}{1 - 2\zeta t}. \quad (40)$$

We can also introduce a velocity with which a fixed space scale factor “travels” along the x^{11} direction:

$$v_{sf} = -\frac{\zeta}{\bar{\alpha}}. \quad (41)$$

The metric (39) describes an initial singularity (space scale factor vanishes) for

$$\bar{\alpha}x^{11} + \zeta t = -1 \quad (42)$$

so the singularity at the other wall ($x^{11} = \pi\rho$) is earlier than at our wall ($x^{11} = 0$). On physical grounds the signal about the beginning of expansion from a point on x^{11} should not reach a nearby point before the expansion has started there.

Therefore the speed of light $|v_{11}|$ should not be greater than $|v_{sf}|$ at each point on x^{11} axis. This condition gives

$$\zeta \geq \bar{\alpha} \sqrt{\frac{1 + \bar{\alpha} x^{11}}{3 + 2\bar{\alpha} x^{11}}} \geq \frac{\bar{\alpha}}{\sqrt{3}}. \quad (43)$$

In terms of our original fields it can be translated as

$$\int_{X^6} G_{t11AB} \omega^{AB} > \frac{4}{\sqrt{3}} \int_{X^6} G_{ABCD} \omega^{AB} \omega^{CD}. \quad (44)$$

This lower bound on $\int_{X^6} G_{t11AB} \omega^{AB}$ is quite surprising. It shows that whenever we switch on nonzero vev for $G_{ab\bar{a}\bar{b}}$ (what we anyway should do due to sources on the walls in the Bianchi identities) while keeping supersymmetry then the solution is physically unstable unless we switch on also sufficiently large vev for $G_{t11a\bar{b}}$. Then the solution is necessarily time-dependent. One should remember however, that the bound can be modified by presently unknown higher order (in $\kappa^{2/3}$) terms.

In conclusion, we have presented the first example of the time dependent solution in the heterotic M-theory that preserves $N = 1$ $d = 4$ supersymmetry. It describes expanding 4-dimensional space-time with shrinking orientifold interval and static Calabi-Yau internal space. As a result the gauge coupling constants do not depend on time while the Newton's constant increases with time. It remains to be seen what are the global properties of such cosmological solutions in M-theory and how they can be used to describe the evolution of the early Universe.

Acknowledgements

K.A.M. was partially supported by the Polish KBN grant 2P03B 03715 (1998-2000) and by the Polish-French Cooperation Grant POLONIUM 1999. M.O. was partially supported by the Polish grant KBN 2 P03B 052 16 (1999-2000).

References

- [1] P. Hořava and E. Witten, Nucl. Phys. B460 (1996) 506; B475 (1996) 94.
- [2] E. Witten, Nucl. Phys. B471 (1996) 135.
- [3] T. Banks and M. Dine, Nucl. Phys. B479 (1996) 173.
- [4] H.P. Nilles, M. Olechowski and M. Yamaguchi, Phys. Lett. B415 (1997) 24; Nucl. Phys. B530 (1998) 43.
- [5] A. Lukas, B.A. Ovrut, and D. Waldram, Nucl. Phys. B495 (1997) 365; Nucl. Phys. B509 (1998) 169; Phys. Rev. D60:86001 (1999); A. Lukas, B.A. Ovrut, Phys. Lett. B437 (1998) 291; H.S. Reall, Phys. Rev. D59:103506 (1999).
- [6] A. Lukas, B.A. Ovrut, K.S. Stelle and D. Waldram, Phys. Rev. D59:86001 (1999).
- [7] A. Lukas, B.A. Ovrut, K.S. Stelle and D. Waldram, Nucl. Phys. B552 (1999) 246; J. Ellis, Z. Lalak, S. Pokorski and S. Thomas, hep-th/9906148.
- [8] K. Benakli, Int. J. Mod. Phys. D8 (1999) 153; Phys. Lett. B447 (1999) 51.
- [9] M. Bremer, M.J. Duff, H. Lu, C.N. Pope, and K.S. Stelle, Nucl. Phys. B543 (1999) 321; S.W. Hawking and H.S. Reall, Phys. Rev. D59:023502 (1999).
- [10] H. Lu, J. Maharana, S. Mukherji, and C.N. Pope, Phys. Rev. D57 (1998) 2219; H. Lu, S. Mukherji, C. N. Pope, and K.-W. Xu, Phys. Rev. D55 (1997) 7926; H. Lu, S. Mukherji, and C.N. Pope, hep-th/9612224; N. Kaloper, I.I. Kogan, and K.A. Olive, Phys. Rev. D57 (1998) 7340; Erratum, ibid. D60:049901 (1999). A. Feinstein and M.A. Vazquez-Mozo, hep-th/9906006; A. Billyard, A. Coley, J. Lidsey and U. Nillson, hep-th/9908102.
- [11] M.J. Duff, P. Hoxha, H. Lu, R.R. Martinez-Acosta, and C.N. Pope, Phys. Lett. B451 (1999) 38; .
- [12] K.A. Meissner, H.P. Nilles and M. Olechowski, hep-th/9905139, to appear in Nucl. Phys. B.